

Boston University School of Law

Scholarly Commons at Boston University School of Law

Faculty Scholarship

2017

Decreasing Smoking but Increasing Stigma? Anti-tobacco Campaigns, Public Health, and Cancer Care

Michael R. Ulrich
Boston University

Follow this and additional works at: https://scholarship.law.bu.edu/faculty_scholarship



Part of the [Health Law and Policy Commons](#)

Recommended Citation

Michael R. Ulrich, *Decreasing Smoking but Increasing Stigma? Anti-tobacco Campaigns, Public Health, and Cancer Care*, 19 *American Medical Association Journal of Ethic* 475 (2017).

Available at: https://scholarship.law.bu.edu/faculty_scholarship/1199

This Article is brought to you for free and open access by Scholarly Commons at Boston University School of Law. It has been accepted for inclusion in Faculty Scholarship by an authorized administrator of Scholarly Commons at Boston University School of Law. For more information, please contact lawlessa@bu.edu.



MEDICINE AND SOCIETY

Decreasing Smoking but Increasing Stigma? Anti-tobacco Campaigns, Public Health, and Cancer Care

Kristen E. Riley, PhD, Michael R. Ulrich, JD, MPH, Heidi A. Hamann, PhD, and Jamie S. Ostroff, PhD

Abstract

Public health researchers, mental health clinicians, philosophers, and medical ethicists have questioned whether the public health benefits of large-scale anti-tobacco campaigns are justified in light of the potential for exacerbating stigma toward patients diagnosed with lung cancer. Although there is strong evidence for the public health benefits of anti-tobacco campaigns, there is a growing appreciation for the need to better attend to the unintended consequence of lung cancer stigma. We argue that there is an ethical burden for creators of public health campaigns to consider lung cancer stigma in the development and dissemination of hard-hitting anti-tobacco campaigns. We also contend that health care professionals have an ethical responsibility to try to mitigate stigmatizing messages of public health campaigns with empathic patient-clinician communication during clinical encounters.

Introduction

Tobacco use remains the leading cause of preventable death in the United States, with cigarette smoking killing more than 480,000 Americans every year [1]. An estimated 41,000 of these deaths among adults are attributable to secondhand smoke exposure [1]. Every day in the US more than 3,800 youths under the age of 18 smoke their first cigarette; an estimated 26 percent of these will become adult smokers [2].

Given the well-established health [consequences of smoking](#), the public health community has established and maintained a comprehensive tobacco control effort, including restrictions on smoking in worksites and other public places, increased tobacco taxation, increased access to evidence-based tobacco treatment, and public health national media campaigns [3]. Collectively, this comprehensive tobacco control effort represents one of the leading public health success stories. In the 50 years since the 1964 Surgeon General's report, *Smoking and Health*, US adult smoking rates have fallen from 43 percent to 18 percent [4].

Although what we'll call "hard-hitting" anti-tobacco public health campaigns—those with fear-arousing messages—have been shown to be the most effective type of anti-tobacco mass-reach health communication interventions, they might have the unintended consequence of stigmatizing those with smoking-related illnesses [5, 6]. In this paper, we explore the ethical dilemma whereby these campaigns are seen as helpful for public health in promoting smoking prevention and cessation but also potentially harmful for persons suffering from tobacco-related illnesses, including lung cancer. We discuss types of stigma and ethical implications, drawing upon concepts such as *respect for persons*. We then make recommendations for public health campaigns to incorporate counter-stigmatizing themes and for health care professionals to use empathic communication to mitigate the effects of stigma on patients with tobacco-related diseases. Finally, we provide direction for future research.

Hard-Hitting Anti-tobacco Public Health Campaigns Are Effective in Reducing Smoking Prevalence

Hard-hitting media anti-smoking campaigns often focus on both raising awareness about the health consequences of smoking and denormalizing smoking behavior, thereby motivating prevention among the general public and motivating smokers specifically toward cessation [7-9]. The term "hard-hitting" has been used to describe ad campaigns that are uncompromisingly direct, often with strong fear-arousing messages and personal stories about negative health consequences of smoking. These types of ads are supported by well-established theories of health behavior change (e.g., the Health Belief Model [10], the theory of planned behavior [11, 12]) that focus broadly on cognitive, emotional, and social processes (e.g., perceived susceptibility to disease, health beliefs regarding the consequences of behavior change, self-efficacy, and social norms) that predict behavior change.

Hard-hitting ads have been shown to be more effective than humorous or neutral educational communication messages at reducing smoking [13]. Most recently, the Tips From Former Smokers™ campaign [14], featuring real people suffering from serious medical conditions as a result of smoking and exposure to secondhand smoke, has been credited with an estimated 1.64 million American smokers making a quit attempt; 100,000 of these smokers are expected to maintain smoking abstinence [8]. Public health leaders assert that the hard-hitting ads are justified by the benefits observed in reducing smoking and related health consequences [5, 15-17]. Although some hard-hitting anti-tobacco campaigns (e.g., graphic warnings on cigarette packs) have been challenged by the tobacco industry [18, 19], the Family Smoking Prevention and Tobacco Control Act of 2009 gives the FDA authority to regulate the tobacco industry [20]. Regardless of these legal challenges, hard-hitting anti-tobacco public health campaigns remain best practice for mass-reach public health communications.

Do Hard-Hitting Anti-tobacco Ad Campaigns Contribute to Stigma?

There are several types of stigma that might be experienced by patients diagnosed with lung cancer: (1) anticipated stigma, or the expectation of discrimination, stereotyping, or prejudice; (2) enacted stigma, which involves actually experiencing discrimination, stereotyping, or prejudice; and (3) internalized stigma, which refers to people's self-endorsing negative feelings and beliefs about themselves [21]. While effective in decreasing smoking rates, hard-hitting anti-tobacco public health campaigns might increase the third kind of stigma. That is, internalization of stigma can result in negative self-appraisal and self-devaluation among persons diagnosed with lung cancer and other tobacco-related diseases [5, 6]. The majority of persons diagnosed with lung cancer report experiencing stigma, often related to guilt, regret, perceived blame, and other negative beliefs about smoking history [16, 22-24].

Stigma is associated with a number of deleterious psychosocial and medical outcomes in lung cancer patients, including delayed diagnoses [25-27], poor quality of life [26], and poor patient-clinician communication [28]. Although there has been limited investigation of stigma and long-term outcomes, stigma may have clear downstream effects, such as reduced treatment adherence and heightened psychosocial distress [24, 28]. One survey found that physicians were more likely to refer breast cancer patients than lung cancer patients for further therapy [29], which could be due to lung cancer stigma—the ubiquitous and damaging nature of which is well established [24, 28, 29].

Previous research has additionally pointed to differential rates of stigma experienced by lung cancer patients who used to or who currently smoke and those who have never smoked. Namely, lung cancer patients who have smoked and those who currently smoke report higher levels of stigma than those who have never smoked [26], although lung cancer patients who have never smoked also report experiencing stigma [26]. Given the epidemiology of lung cancer, health care professionals might assume that a patient's lung disease is acquired "firsthand" as opposed to "secondhand" or without smoking exposure at all. As stigma is experienced by patients across this continuum of smoking exposures, the salience of this ethical debate is relevant for current, former, and never smokers—all those suffering from illnesses associated with smoking.

An Ethical Dilemma

While recognizing that the public health goals of tobacco prevention and cessation remain paramount, an ethical question arises as to whether these ads should continue to be hard-hitting or whether public health communication messages should be reframed to try to reduce stigma and blame that could be experienced by the 16 million Americans living with smoking-related diseases [30]. Looking at denormalization of smoking through a purely utilitarian lens renders a favorable assessment, as evidenced by a 12 percent drop in the smoking rate of 18- to 29-year-olds in the US from 2005 to 2015 [31]. However, when viewing hard-hitting anti-tobacco public health campaigns as

sanctioned social stigmatization in the context of people suffering from nicotine addiction and related medical illnesses, the “benefits” of these anti-tobacco ads should be tempered [32]. Internalized stigma (e.g., self-blame, shame, or guilt) could result in low self-esteem as people question their identity and self-worth. In its extreme form, stigma can be thought to “turn the individual into his own jailor, his own chorus of denunciation” [33].

Mental health clinicians caring for the psychosocial needs of cancer patients and health care ethicists have questioned whether the public health benefits are worth the “incidental” costs of stigma for individual patients [16, 34, 35]. Some health scientists have labeled anti-tobacco public health campaigns “demoralizing” [22] and “victim blaming” [6]. Additionally, hard-hitting campaigns could extend lung cancer stigma to any person who suffers from any smoking-related illness, regardless of the patient’s actual smoking history [24]. This “guilt by association” can be especially difficult for those with secondhand or even no prior tobacco exposure who perceive others’ negative attitudes as based on false assumptions about the nature and scope of their disease culpability. Given the current demographics of tobacco use, these campaigns might further stigmatize low-income and other vulnerable populations of smokers, who currently represent the majority of tobacco users [22]. And people who already feel disempowered tend to feel even more resentful, defensive, and demoralized after exposure to anti-tobacco campaigns [17, 36]. As a result, hard-hitting anti-tobacco ads could exacerbate health disparities and discourage access to high-quality health care.

An important ethical question is how much iatrogenic stigma should matter if hard-hitting campaigns are successful in preventing tobacco use and motivating smoking cessation as public health goals. Stigma and associated distress certainly matter at a level of clinicians interacting with individual patients diagnosed with lung cancer or other tobacco-related diseases. How much should an individual’s experience of stigma matter at a macro- and public health level of disease prevention? If the overarching goal is to reduce the negative health effects of tobacco use and smoking, whether the result of firsthand or secondhand use, the potential stigmatizing impact of anti-tobacco ads on those who are already suffering from tobacco-related illnesses such as lung cancer cannot be ignored.

Stigma is not benign and has been shown to be associated with lung cancer patients’ avoidance or delay of seeking medical care [25], resulting in downstream risk of worsening lung cancer morbidity and mortality. While public health principles often emphasize prevention, stigma does not exclude those populations that prevention efforts have failed to reach. Meanwhile, the ethical principle of respect for persons and appreciating the intrinsic value of each individual requires that those who are suffering from tobacco-related illnesses, such as lung cancer, be treated with equity and justice. Health care professionals taking their ethical obligation of nonmaleficence seriously

should certainly be concerned about their roles in whether and how their individual patients experience stigma as a result of their specific actions or communications.

What Should Be Done?

Because anti-tobacco public health campaigns have been effective in reducing population smoking rates, banning hard-hitting ads completely would be shortsighted. Our attempt to raise awareness about the impact of lung cancer stigma is not to suggest that public health campaigns refrain from educating the public about the unquestionable, far-reaching health hazards of smoking. Rather, we offer several recommendations for addressing the iatrogenic consequences of hard-hitting anti-tobacco campaigns.

First, public health campaigns could highlight counter-stigma themes. One such theme is the unscrupulous, predatory nature of big tobacco as an industry. Emphasizing how much money is spent annually by the tobacco industry on tobacco advertising and social marketing has been a compelling theme for prior anti-tobacco campaigns, particularly those targeting prevention of youth smoking [37-39]. The Lung Cancer Alliance's campaign, "No One Deserves to Die of Lung Cancer," serves as an excellent example of an effective public health campaign that acknowledges the dangerous nature of cigarette smoking while also emphasizing compassion and a nonjudgmental stance by using the ironic message that certain segments of the population (e.g., cat ladies, hipsters) deserve to die [40]. Ads that provide self-affirming messages (e.g., the value of raising a family or maintaining health) might buffer against defensive processing—dismissal of a health message perceived as personally threatening—because it has been shown that self-affirmation prior to exposure to graphic images on cigarette pack warnings reduces such defensive processing [41]. Recent research shows that gain-framed messages—those that highlight benefits of quitting rather than costs of smoking—might be more effective for smokers who feel helpless and demoralized in their quitting efforts [42]. We also recommend ads that encourage the use of evidence-based smoking behavior change strategies and promote self-efficacy in quitting. Finally, given that lung cancer stigma can intersect with social and structural hierarchies such as power, culture, and privilege [43], it would seem important for public health campaigns to target all tobacco users, not just ethnic minorities and tobacco users of low socioeconomic status [32].

Second, health care professionals treating patients with lung cancer can communicate empathically to build patients' resilience and try to help inoculate them to the stigmatizing effects of anti-tobacco health campaigns [6, 24, 26, 28]. One study found that physicians miss 90 percent of opportunities for demonstrating empathy in lung cancer care [44]. Physicians have noted the challenge of advising their patients to quit smoking while concurrently managing patients' emotional distress following cancer diagnosis and treatment [6]. Good patient-clinician communication has been associated with lower levels of stigma in the health care setting [28]. Building resilience in lung cancer patients and those with tobacco-related illnesses through empathic responses

and problem-focused strategies may mitigate the negative consequences of stigma resulting from hard-hitting anti-tobacco campaigns [45]. We currently are working to develop and evaluate an empathic, nonjudgmental communication skills training module for health care professionals treating patients with lung cancer that focuses on taking a detailed tobacco history, advising current smokers to quit, and making a reliable referral for tobacco treatment services.

Additional research is needed to determine how anti-tobacco campaigns can minimize the internalized stigma of patients living with tobacco-related diseases without compromising the campaigns' strong public health effectiveness. For example, public health campaigns are often pretested using focus groups; new candidate ads could be assessed for whether and to what extent they generate stigma and unintended consequences such as shame and guilt. To our knowledge, the Tips campaign has not examined whether patients with lung and other tobacco-related conditions experience heightened stigma and regret. We recommend eliciting patient perspectives early in the development of anti-tobacco campaigns. There is much to be learned from other public health campaigns grappling with similar concerns (e.g., risky sexual and drug use behaviors and HIV/AIDS, alcohol and driving, obesity, and sun exposure). The Joint United Nations Programme on HIV/AIDS (UNAIDS) has suggested that negative views of [people living with HIV](#) can be attributed largely to stigma and ignorance about the harm of stigma and moral judgment, which is likely germane to those suffering from tobacco-related diseases [46]. Accordingly, the HIV/AIDS public health community has made a concerted effort to examine the impact of stigma and embark on multipronged efforts to counter stigma with educational programs targeting specific vulnerable populations, in addition to addressing the role of health care professionals in exacerbating the effects of stigma [47].

Conclusion

Overall, hard-hitting anti-tobacco public health campaigns work, although they might also inadvertently increase stigma among lung cancer patients, leading to deleterious downstream psychosocial and medical outcomes for this vulnerable population. Specific recommendations include shifting the focus of public health campaigns away from patient blaming and emphasizing clinician-level empathic communication interventions. Further research and attention are needed to ensure that hard-hitting anti-tobacco campaigns find the "sweet spot" for maximizing tobacco control while minimizing stigma experienced by lung cancer patients and those suffering from tobacco-related illnesses. Researchers, leaders of nonprofit organizations, government, hospital systems, health care professionals, and patient advocates can all be involved and accountable for decreasing stigma directed towards lung cancer patients.

References

1. Centers for Disease Control and Prevention. Tobacco-related mortality. https://www.cdc.gov/tobacco/data_statistics/fact_sheets/health_effects/tobacco_related_mortality/. Updated December 1, 2016. Accessed March 10, 2017.
2. US Department of Health and Human Services. *Preventing Tobacco Use Among Youth and Young Adults: A Report of the Surgeon General*. Rockville, MD: US Department of Health and Human Services; 2012. <https://www.surgeongeneral.gov/library/reports/preventing-youth-tobacco-use/full-report.pdf>. Accessed March 31, 2017.
3. Centers for Disease Control and Prevention. *Best Practices for Comprehensive Tobacco Control Programs—2014*. Atlanta, GA: Centers for Disease Control and Prevention; 2014. https://www.cdc.gov/tobacco/stateandcommunity/best_practices/pdfs/2014/comprehensive.pdf. Accessed March 10, 2017.
4. US Department of Health and Human Services. *The Health Consequences of Smoking—50 Years of Progress: A Report of the Surgeon General*. Atlanta, GA: Centers for Disease Control and Prevention; 2014. <http://www.surgeongeneral.gov/library/reports/50-years-of-progress/full-report.pdf>. Accessed September 15, 2016.
5. Bayer R. Stigma and the ethics of public health: not can we but should we. *Soc Sci Med*. 2008;67(3):463-472.
6. Chapple A, Ziebland S, McPherson A. Stigma, shame, and blame experienced by patients with lung cancer: qualitative study. *BMJ*. 2004;328(7454):1470. <http://www.bmj.com/content/328/7454/1470>. Accessed March 30, 2017.
7. Duke JC, Hansen H, Kim AE, Curry L, Allen J. The use of social media by state tobacco control programs to promote smoking cessation: a cross-sectional study. *J Med Internet Res*. 2014;16(7):e169.
8. McAfee T, Babb S, McNabb S, Fiore MC. Helping smokers quit—opportunities created by the Affordable Care Act. *N Engl J Med*. 2015;372(1):5-7.
9. Johnson KA, Stewart S, Rosenfield D, Steeves D, Zvolensky MJ. Prospective evaluation of the effects of anxiety sensitivity and state anxiety in predicting acute nicotine withdrawal symptoms during smoking cessation. *Psychol Addict Behav*. 2012;26(2):289-297.
10. Rosenstock IM. The health belief model and preventive health behavior. *Health Educ Behav*. 1974;2(4):354-386.
11. Ajzen I. From intentions to actions: a theory of planned behavior. In: Kuhl J, Beckmann J. *Action Control: From Cognition to Behavior*. Berlin, Germany: Springer Berlin-Heidelberg; 1985:11-39.
12. Fishbein M, Ajzen I. *Belief, Attitude, Intention, and Behavior: An Introduction to Theory and Research*. Reading, MA: Addison-Wesley; 1975:6.

13. Durkin SJ, Biener L, Wakefield MA. Effects of different types of antismoking ads on reducing disparities in smoking cessation among socioeconomic subgroups. *Am J Public Health*. 2009;99(12):2217-2223.
14. Centers for Disease Control and Prevention. TIPS From Former Smokers™. <https://www.cdc.gov/tobacco/campaign/tips/>. Updated January 19, 2017. Accessed March 13, 2017.
15. Alderman J, Dollar KM, Kozlowski LT. Commentary: understanding the origins of anger, contempt, and disgust in public health policy disputes: applying moral psychology to harm reduction debates. *J Public Health Policy*. 2010;31(1):1-16.
16. Bayer R, Stuber J. Tobacco control, stigma, and public health: rethinking the relations. *Am J Public Health*. 2006;96(1):47-50.
17. Lupton D. The pedagogy of disgust: the ethical, moral and political implications of using disgust in public health campaigns. *Crit Public Health*. 2015;25(1):4-14.
18. Bayer R, Johns D, Colgrove J. The FDA and graphic cigarette-pack warnings—thwarted by the courts. *New Engl J Med*. 2013;369(3):206-208.
19. Mickle T. Antitobacco groups sue FDA to require graphic warning labels on cigarette packs. *Wall Street Journal*. October 4, 2016. <https://www.wsj.com/articles/antitobacco-groups-sue-fda-to-require-graphic-warning-labels-on-cigarette-packs-1475599261>. Accessed March 30, 2017.
20. Family Smoking Prevention and Tobacco Control Act of 2009, Pub L No. 111-31, 123 Stat 1776. <https://www.gpo.gov/fdsys/pkg/PLAW-111publ31/html/PLAW-111publ31.htm>. Accessed March 30, 2017.
21. Sweeney SM, Venable PA. The association of HIV-related stigma to HIV medication adherence: a systematic review and synthesis of the literature. *AIDS Behav*. 2016;20(1):29-50.
22. Bell K, Salmon A, Bowers M, Bell J, McCullough L. Smoking, stigma and tobacco “denormalization”: further reflections on the use of stigma as a public health tool. A commentary on *Social Science & Medicine's* stigma, prejudice, discrimination and health special issue (67:3). *Soc Sci Med*. 2010;70(6):795-799.
23. Gritz ER, Vidrine DJ, Fingeret MC. Smoking cessation: a critical component of medical management in chronic disease populations. *Am J Prev Med*. 2007;33(suppl 6):S414-S422.
24. Hamann HA, Ostroff JS, Marks EG, Gerber DE, Schiller JH, Lee SJ. Stigma among patients with lung cancer: a patient reported measurement model. *Psychooncology*. 2014;23(1):81-92.
25. Carter-Harris L, Hermann CP, Schreiber J, Weaver MT, Rawl SM. Lung cancer stigma predicts timing of medical help-seeking behavior. *Oncol Nurs Forum*. 2014;41(3):e203-e210.
26. Cataldo JK, Jahan TM, Pongquan VL. Lung cancer stigma, depression, and quality of life among ever and never smokers. *Eur J Oncol Nurs*. 2012;16(3):264-269.
27. Chambers SK, Dunn J, Occhipinti S, et al. A systematic review of the impact of stigma and nihilism on lung cancer outcomes. *BMC Cancer*. 2012;12:184.

<https://bmccancer.biomedcentral.com/articles/10.1186/1471-2407-12-184>. Accessed March 30, 2017.

28. Shen MJ, Hamann HA, Thomas AJ, Ostroff JS. Association between patient-provider communication and lung cancer stigma. *Support Care Cancer*. 2016;24(5):2093-2099.
29. Wassenaar TR, Eickhoff JC, Jarzemsky DR, Smith SS, Larson ML, Schiller JH. Differences in primary care clinicians' approach to non-small cell lung cancer patients compared with breast cancer. *J Thorac Oncol*. 2007;2(8):722-728.
30. Centers for Disease Control and Prevention. Fast facts: diseases and death. https://www.cdc.gov/tobacco/data_statistics/fact_sheets/fast_facts/. Updated December 20, 2016. Accessed March 12, 2017.
31. Nader N, Liu D. In US, young adults' cigarette use is down sharply. *Gallup*. December 10, 2015. <http://www.gallup.com/poll/187592/young-adults-cigarette-down-sharply.aspx>. Accessed March 12, 2017.
32. Burris S. Disease stigma in public health law. *J Law Med Ethics*. 2002;30(2):179-190.
33. Burris S. Stigma, ethics and policy: a commentary on Bayer's "Stigma and the ethics of public health: not can we but should we." *Soc Sci Med*. 2008;67(3):476.
34. Guttman N, Salmon CT. Guilt, fear, stigma and knowledge gaps: ethical issues in public health communication interventions. *Bioethics*. 2004;18(6):531-552.
35. Fairchild AL, Bayer R, Colgrove J. The renormalization of smoking? E-cigarettes and the tobacco "endgame." *N Engl J Med*. 2014;370(4):293-295.
36. Hastings G, Stead M, Webb J. Fear appeals in social marketing: strategic and ethical reasons for concern. *Psychol Mark*. 2004;21(11):961-986.
37. Farrelly MC, Davis KC, Duke J, Messeri P. Sustaining "truth": changes in youth tobacco attitudes and smoking intentions after 3 years of a national antismoking campaign. *Health Educ Res*. 2009;24(1):42-48.
38. Farrelly MC, Davis KC, Haviland ML, Messeri P, Healton CG. Evidence of a dose-response relationship between "truth" antismoking ads and youth smoking prevalence. *Am J Public Health*. 2005;95(3):425-431.
39. Malone RE, Grundy Q, Bero LA. Tobacco industry denormalisation as a tobacco control intervention: a review. *Tob Control*. 2012;21(2):162-170.
40. Lung Cancer Alliance. No one deserves to die of lung cancer. <http://www.noonedeservestodie.org/>. Accessed March 12, 2017.
41. Harris PR, Mayle K, Mabbott L, Napper L. Self-affirmation reduces smokers' defensiveness to graphic on-pack cigarette warning labels. *Health Psychol*. 2007;26(4):437-446.
42. Mead EL. *Moving Beyond Fear: Exploring Perceptions of Theory-Based Graphic Warning Labels Among Low-Income, Urban Smokers* [dissertation]. Baltimore, MD: Johns Hopkins University; 2014. <https://jscholarship.library.jhu.edu/bitstream/handle/1774.2/39316/MEAD-DISSERTATION-2014.pdf?sequence=1&isAllowed=y>. Accessed March 31, 2017.

43. Knight J. Updated definition of internationalization. *Int High Educ.* 2015;(33):2-3.
44. Morse DS, Edwardsen EA, Gordon HS. Missed opportunities for interval empathy in lung cancer communication. *Arch Intern Med.* 2008;168(17):1853-1858.
45. Thompson L, Barnett JR, Pearce JR. Scared straight? Fear-appeal anti-smoking campaigns, risk, self-efficacy and addiction. *Health Risk Soc.* 2009;11(2):181-196.
46. Brent RJ. The value of reducing HIV stigma. *Soc Sci Med.* 2016;151:233-240.
47. Pellowski JA, Kalichman SC, Matthews KA, Adler N. A pandemic of the poor: social disadvantage and the US HIV epidemic. *Am Psychol.* 2013;68(4):197-209.

Kristen E. Riley, PhD, is a postdoctoral research fellow at Memorial Sloan Kettering Cancer Center in New York City, where she studies health behavior decision making and tobacco cessation for cancer prevention.

Michael R. Ulrich, JD, MPH, is an assistant professor in the Center for Health Law, Ethics & Human Rights and the Department of Health, Law, and Policy Management at Boston University School of Public Health. He studies public health, ethics, and law.

Heidi A. Hamann, PhD, is an associate professor in the Departments of Psychology and Family and Community Medicine at the University of Arizona in Tucson. Her primary research focuses on psychosocial and behavioral concerns, including stigma of lung cancer patients and survivors.

Jamie S. Ostroff, PhD, is the chief of the behavioral sciences service in the Department of Psychiatry and Behavioral Sciences at Memorial Sloan Kettering Cancer Center and a professor of psychology in the Department of Healthcare Policy and Research at Weill Cornell Medical College, both in New York City. Her cancer prevention and control research focuses on tobacco cessation and stigma experienced by patients with lung cancer.

Acknowledgements

This work was supported in part through a cancer center support grant (P30CA0087748) from the National Cancer Institute to the Memorial Sloan Kettering Cancer Center, a training grant from the National Cancer Institute (T32CA009461) (KER), the National Cancer Institute (R03CA154016) and the National Lung Cancer Partnership and its North Carolina Chapter (Young Investigator Award, HAH), and the Lung Cancer Research Foundation (JSO). We would like to thank Jack Burkhalter, PhD, Lisa Carter-Harris, PhD, Maureen Rigney, LCSW, and Christine Sheffer, PhD, for their constructive feedback on an earlier version of this paper.

Related in the *AMA Journal of Ethics*

[The Ethics of Requiring Employees To Quit Smoking](#), January 2007

[Health Effects of Smoking and the Benefits of Quitting](#), January 2011

[HIV Stigma and Discrimination Persist, Even in Health Care](#), December 2009
[Teaching Smoking Cessation in U.S. Medical Schools: A Long Way to Go](#), January 2007
[Use of Images in Public Health Campaigns](#), August 2007

The viewpoints expressed in this article are those of the author(s) and do not necessarily reflect the views and policies of the AMA.

**Copyright 2017 American Medical Association. All rights reserved.
ISSN 2376-6980**